



Predation by a Brown Rainbow Boa (*Epicrates maurus*) on a Khaki Ameiva (*Ameiva bifrontata*) in Colombia

Freddy A. Grisales-Martínez^{1,2}, M. Silvana Cárdenas-Ortega³, and Beatriz E. Rendón-Valencia¹

¹Servicios Ambientales y Geográficos S.A., Calle 11B No. 40A-130 Piso 4, Medellín, Colombia (fgrisales@sag-sa.com; beatriz.rendon@sag-sa.com)

²Grupo Herpetológico de Antioquia (GHA), Instituto de Biología, Universidad de Antioquia, Calle 67 # 53-108, Bloque 7-121, A.A. 1226, Medellín, Colombia (griso88@gmail.com [corresponding author])

³Grupo de Ecología y Diversidad de Anfibios y Reptiles (GEDAR), Facultad de Ciencias Exactas y Naturales, Universidad de Caldas, Calle 65 # 26-10, A. A. 275, Manizales, Colombia (silvana9420@gmail.com)

The Brown Rainbow Boa (*Epicrates maurus* Gray 1849), a non-venomous constrictor that can grow to 1.5 m SVL, is nocturnally active, terrestrial or semi-arboreal, and occurs in a variety of habitats, including dry forests, savannas, and open areas, even those with some level of anthropogenic disturbance (Pérez-Santos and Moreno 1988; Savage 2002; Wallach et al. 2014; Rojas-Murcia et al. 2016). This species is widely distributed at elevations to 1,200 m in central and northern South America (Savage 2002; Uetz et al. 2024).

Little is known about the prey of these snakes, with reports limited to a Neotropical Ameiva (*Ameiva ameiva*) (Starace 2013), a Greater Sac-winged Bat (*Saccopteryx bilineata*) (Aya-Cuero et al. 2019), a European Rabbit (*Oryctolagus cuniculus*) (Salcedo-Rivera et al. 2021), and a Short-tailed Cane Mouse (*Zygodontomys brevicauda*) (Salazar-Guzmán et al. 2022). Also, filial cannibalism of dead hatchlings and undeveloped eggs was documented by Lourdaís et al. (2005).

At 1520 h on 11 July 2024, we captured a juvenile female *E. maurus* (SVL 635 mm, TL 87 mm, mass 114 g) in a pile of wood and soil in tropical dry forest in the Upper Magdalena River Basin, Colombia, during rescue and deterrence activities for the “Puerta de Oro” solar project located in a rural area of the Municipality of Guaduas, Department of Cundinamarca, Colombia (5.021667, -74.726889; elev. 227 m asl). After capture, we placed the boa in a bucket in which, five minutes later, it regurgitated an adult male Khaki Ameiva (*Ameiva bifrontata* Cope 1862) (SVL 125 mm, TL 272 mm, mass 57 g) (Fig. 1). The lizard, which was intact and showed no signs of digestion, had probably been ingested shortly before the snake was captured. The snake was released in the project’s faunal receptor area; the lizard was preserved and deposited in the Museo de Herpetología de la Universidad de Antioquia (MHUA MHUA-R 16022).

The ingested lizard had half the mass of the boa. Size relationships between predator and prey in snakes are often highly disproportionate (e.g., Glaudas et al. 2018). In particular, for boids, the predator-prey size relationship shows that relative decrease in head length with increasing body size, allows smaller snakes to exploit relatively larger prey than adults (Pizzatto et al. 2009).

To the best of our knowledge, this is only the second report of any snake in the genus *Epicrates* preying on an ameiva. Ameivas (Family Teiidae) are diurnally active and very agile, whereas *E. maurus* is mainly nocturnal, with foraging and feeding occurring primarily during evening



Figure 1. A Brown Rainbow Boa (*Epicrates maurus*) and a Khaki Ameiva (*Ameiva bifrontata*) in the bucket where the snake’s prey was regurgitated. Photograph by Freddy Alexander Grisales-Martínez.

hours, while during the day it typically shelters under cover (Kissoon 2011). Consequently, most of the documented prey species are small mammals that exhibit crepuscular and/or nocturnal behavior (Aya-Cuero et al. 2019; Salcedo-Rivera et al. 2021; Salazar-Guzmán et al. 2022). However, at least some boids feed on both diurnal and nocturnal prey while employing a combination of active searching and sit-and-wait foraging strategies (Henderson and Pauers 2012; Rodríguez-Cabrera et al. 2020). Furthermore, as in this case and the one previous report of *E. maurus* consuming an ameiva (Starace 2013), juveniles of at least some boids tend to consume ectothermic prey, whereas adults are more likely to prey on endotherms (Pizzatto et al. 2009; Henderson and Pauers 2012; Henderson et al. 2021).

Acknowledgements

We thank Puerta de Oro for allowing us to use the data from the project's fauna rescue and deterrence activities for this publication; and the biologists, veterinarians, and field assistants who contributed to the project. The first author thanks the Museo de Herpetología de la Universidad de Antioquia (MHUA) for providing workspace.

Literature Cited

- Aya-Cuero, C.A., C.H. Cáceres-Martínez, and D.A. Esquivel. 2019. First record of predation on Greater sac-winged bat, *Saccopteryx bilineata* (Chiroptera: Emballonuridae), by the Colombian rainbow boa, *Epicrates maurus* (Serpentes: Boidae). *Herpetology Notes* 12: 815–817.
- Glaudas, X., K.L. Glennon, M. Martins, L. Luiselli, S. Fearn, D.F. Trembath, D. Jelić, and G.J. Alexander. 2018. Foraging mode, relative prey size and diet breadth: A phylogenetically explicit analysis of snake feeding ecology. *Journal of Animal Ecology* 88: 757–767. <https://doi.org/10.1111/1365-2656.12972>.
- Henderson, R.W. and M.J. Pauers. 2012. On the diets of Neotropical treeboas (Squamata: Boidae: *Corallus*). *South American Journal of Herpetology* 7: 172–180. <http://dx.doi.org/10.2994/057.007.0207>.
- Henderson, R.W., E.T. Hileman, R.A. Sajdak, B.C. Harrison, R. Powell, and D.R. Bradke. 2021. Effects of body size, diet, and transience on the demography of the arboreal boid snake *Corallus grenadensis* on Carriacou (Grenada Grenadines, West Indies). *Population Ecology* 63: 177–188. <https://doi.org/10.1002/1438-390X.12079>.
- Kissoon, S. 2011. *Epicrates maurus* (Rainbow Boa or Velvet Mapepire). *The Online Guide to the Animals of Trinidad and Tobago*. <https://sta.uwi.edu/fst/lifesciences/sites/default/files/lifesciences/documents/ogatt/Eira_barbara%20-%20Tayra.pdf>.
- Lourdais, O., F. Brischoux, R. Shine, and X. Bonnet. 2005. Adaptive maternal cannibalism in snakes (*Epicrates cenchria maurus*, Boidae). *Biological Journal of the Linnean Society* 84: 767–774. <https://doi.org/10.1111/j.1095-8312.2005.00436.x>.
- Pérez-Santos, C. and A.G. Moreno 1988. *Ofidios de Colombia*. Museo Regionale di Scienze Naturali, Torino, Italy.
- Pizzatto, L., O. Marques, and K. Facure. 2009. Food habits of Brazilian boid snakes: overview and new data, with special reference to *Corallus hortulanus*. *Amphibia-Reptilia* 30: 533–544. <https://doi.org/10.1163/156853809789647121>.
- Rodríguez-Robles, J.A. 2002. Feeding ecology of North American gopher snakes (*Pituophis catenifer*, Colubridae). *Biological Journal of the Linnean Society* 77: 165–183. <https://doi.org/10.1046/j.1095-8312.2002.00098.x>.
- Rojas Murcia, L.E., J.E. Carvajal Cogollo, and J.A. Cabrejo Bello. 2016. Reptiles del bosque seco estacional en el Caribe Colombiano: Distribución de los hábitats y del recurso alimentario. Reptiles from the seasonal dry forest the Caribbean Region: Distribution of habitat and use of food resource. *Acta Biologica Colombiana* 21: 365–377. <https://doi.org/http://dx.doi.org/10.15446/abc.v21n2.49393>.
- Salazar-Guzmán, A.M., C. Cárdenas-González, A. Niño-Reyes, A.J. Lozano-Flórez, and J.A. Rojas-Morales. 2022. Nueva presa para la boa arcoíris colombiana, *Epicrates maurus* (Gray, 1849) (Squamata, Serpentes, Boidae): el ratón de caña de cola corta *Zygodontomys brevicauda* (J.A. Allen & F.M. Chapman, 1893). *Revista Latinoamericana de Herpetología* 5: 69–73. <https://doi.org/10.22201/fc.25942158e.2022.3.489>.
- Salcedo-Rivera, G.A., J. Vanegas-Arroyo, C.J. Castillo, J.A. Díaz-Pérez, and D. Montes-Vergara. 2021. On the human-snake conflict: Predation attempt of *Oryzomys cuniculus* (Linnaeus, 1758) by *Epicrates maurus* Gray, 1849 in a farm in the Caribbean Region of Colombia. *Herpetology Notes* 14: 941–943.
- Savage, J.M. 2002. *The Amphibians and Reptiles of Costa Rica: A Herpetofauna Between Two Continents, Between Two Seas*. University of Chicago Press, Chicago, Illinois, USA.
- Starace, F. 2013. *Serpents et amphibiens de Guyane française*. Ibis Rouge Éditions, Matoury, Guyane.
- Uetz, P., P. Freed, R. Aguilar, F. Reyes, J. Kudera, and J. Hošek (eds.). 2024. *The Reptile Database*. <<http://www.reptile-database.org>>.
- Wallach, V., K.L. Williams, and J. Boundy. 2014. *Snakes of the World: A Catalogue of Living and Extinct Species*. CRC Press, Boca Raton, Florida, USA.